Walnutdale Dairy

4-24-13 CNMP Field Additions Records

Prepared by:



April 24, 2013



April 24, 2013

Walnutdale Dairy - CAFO

Field Additions - Spring 2013

Mr. Bruce Washburn,

Mr. (a) (b) (c) is submitting these fields for review and addition to his CNMP for the 2013 crop season. Physical evaluations of slopes and setbacks and a crop plan have been developed and are submitted for review. Soil tests were taken and are currently at A&L Labs. When samples are reported, they will be included with a crop plan and manure application plan reflecting targeted crops and fertility needs.

The fields attached will not receive manure applications when the ground is frozen or snow covered, but MARI evaluations have been included for reference. MARI evaluations will be updated when soil tests are reported.

Manure Application Rates will be established when soil tests are reported.

Crop Plans & Recommendations:

The Home West, East and South fields are alfalfa; the remaining Middle field is going to be planted for grain corn. Recommendations for crop production and manure utilization will be included with the soil test update.

RUSLE2 Evaluations:

All fields are within the tolerable range of predicted soil erosion.

Wind Erosion Evaluations:

Wind erosion reviews were conducted on the fields with I values of 86 or higher. The predicted rates of wind erosion on all fields are below the tolerable limit.

Please review the information attached and contact [0] (6) or myself with any questions on these fields.

Thank you,

Jason Stegink – Green Valley Agricultural, Inc.

Walnutdale Dairy Spreading







Section(s): 14&23 Fields:

Home West, Middle, East and South


North:



| 建筑 | Never apply manure under any conditions (min. 100' witdth) |
|-----------|--|
| | Applications under excellent conditions with 24hr incorp. |

Revised: 4-24-13





MAP LEGEND

Area of Interest (AOI)

A

Area of Interest (AOI)

Soils

Soil Map Units

Special Point Features

Blowout

Borrow Pit

Clay Spot

Closed Depression

Gravel Pit

Gravelly Spot

Landfill

A Lava Flow

Marsh or swamp

Mine or Quarry

Miscellaneous Water

Perennial Water

Rock Outcrop

+ Saline Spot

. Sandy Spot

Severely Eroded Spot

Sinkhole

3 Slide or Slip

g Sodic Spot

Spoil Area

↑ Stony Spot

Wery Stony Spot

Wet Spot

Other

Special Line Features

2

Gully

. . Short Steep Slope

Other

Political Features

0

Cities

Water Features

~

Streams and Canals

Transportation

+++

Rails

-

Interstate Highways

~

US Routes Major Roads



Local Roads

MAP INFORMATION

Map Scale: 1:10,500 if printed on A size (8.5" × 11") sheet.

The soil surveys that comprise your AOI were mapped at 1:15,840.

Please rely on the bar scale on each map sheet for accurate map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL: http://websoilsurvey.nrcs.usda.gov Coordinate System: UTM Zone 16N NAD83

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.


Soil Survey Area: Allegan County, Michigan Survey Area Data: Version 9, Sep 28, 2012

Date(s) aerial images were photographed: 6/22/2005

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

| | Allegan County, Michigan (M | 1005) | |
|---------------------------|---|--------------|----------------|
| Map Unit Symbol | Map Unit Name | Acres in AOI | Percent of AOI |
| 2 | Glendora loamy sand | 14.9 | 2.9% |
| 5 | Houghton muck | 38.6 | 7.6% |
| 10B | Oakville fine sand, 0 to 6 percent slopes | 5.7 | 1.1% |
| 10C | Oakville fine sand, 6 to 18 percent slopes | 36.2 | 7.1% |
| 11B | Oshtemo-Chelsea complex, 0 to 6 percent slopes | 21.2 | 4.2% |
| 14C | Marlette loam, 6 to 12 percent slopes | 25.7 | 5.1% |
| 14D | Marlette loam, 12 to 18 percent slopes | 3.2 | 0.6% |
| 16B | Capac loam, 0 to 6 percent slopes | 70.3 | 13.9% |
| 26A | Pipestone sand, 0 to 4 percent slopes | 17.2 | 3.4% |
| 27B | Metea loamy fine sand, 1 to 6 percent slopes | 125.4 | 24.7% |
| 27C | Metea loamy fine sand, 6 to 12 percent slopes | 20.5 | 4.0% |
| 28A | Rimer loamy sand, 0 to 4 percent slopes | 38.9 | 7.7% |
| 29 | Cohoctah silt loam | 11.8 | 2.3% |
| 30 | Colwood silt loam | 4.1 | 0.8% |
| 39 | Granby loamy sand | 3.4 | 0.7% |
| 42B | Metamora sandy loam, 1 to 4 percent slopes | 1.0 | 0.2% |
| 44C | Chelsea loamy fine sand, 6 to 12 percent slopes | 17.3 | 3.4% |
| 48 | Belleville loamy sand | 2.8 | 0.6% |
| 49A | Tedrow fine sand, 0 to 4 percent slopes | 3.1 | 0.6% |
| 50 | Aquents and Histosols, ponded | 3.5 | 0.7% |
| 51A | Thetford loamy fine sand, 0 to 4 percent slopes | 4.1 | 0.8% |
| 53B | Oakville fine sand, loamy substratum, 0 to 6 percent slopes | 12.7 | 2.5% |
| 75B | Marlette-Capac loams, 1 to 6 percent slopes | 25.5 | 5.0% |
| W | Water | 0.4 | 0.1% |
| Totals for Area of Intere | est | 507.6 | 100.0% |



MAP LEGEND

Area of Interest (AOI)

A

Area of Interest (AOI)

Soils

Soil Map Units

Special Point Features

Blowout

Borrow Pit

Clay Spot

Closed Depression

Gravel Pit

.. Gravelly Spot

A Landfill

∧ Lava Flow

الم

Mine or Quarry

Miscellaneous Water

Perennial Water

Rock Outcrop

+ Saline Spot

Sandy Spot

Severely Eroded Spot

Sinkhole

3 Slide or Slip

g Sodic Spot

Spoil Area

Stony Spot

Very Stony Spot

Wet Spot

Other

Special Line Features

کی و

Gully

. Short Steep Slope

Other

Political Features

Cities

Water Features

Streams and Canals

Transportation

+++ F

+ Raiis

-

Interstate Highways

~

US Routes

~~

Major Roads

Local Roads

MAP INFORMATION

Map Scale: 1:9,180 if printed on A size (8.5" × 11") sheet.

The soil surveys that comprise your AOI were mapped at 1:15,840.

Please rely on the bar scale on each map sheet for accurate map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL: http://websoilsurvey.nrcs.usda.gov

Coordinate System: UTM Zone 16N NAD83

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Allegan County, Michigan Survey Area Data: Version 9, Sep 28, 2012

Date(s) aerial images were photographed: 6/22/2005

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

| Allegan County, Michigan (MI005) | | | | | | | |
|----------------------------------|---|--------------|----------------|--|--|--|--|
| Map Unit Symbol | Map Unit Name | Acres in AOI | Percent of AOI | | | | |
| 2 | Glendora loamy sand | 37.4 | 11.4% | | | | |
| 10B | Oakville fine sand, 0 to 6 percent slopes | 19.6 | 6.0% | | | | |
| 14C | Marlette loam, 6 to 12 percent slopes | 4.2 | 1.3% | | | | |
| 16B | Capac loam, 0 to 6 percent slopes | 91.7 | 27.9% | | | | |
| 26A | Pipestone sand, 0 to 4 percent slopes | 55.7 | 16.9% | | | | |
| 27B | Metea loamy fine sand, 1 to 6 percent slopes | 49.1 | 14.9% | | | | |
| 27C | Metea loamy fine sand, 6 to 12 percent slopes | 13.7 | 4.1% | | | | |
| 28A | Rimer loamy sand, 0 to 4 percent slopes | 7.2 | 2.2% | | | | |
| 30 | Colwood silt loam | 5.7 | 1.7% | | | | |
| 34 | Aquents, sandy and loamy | 0.2 | 0.1% | | | | |
| 39 | Granby loamy sand | 5.6 | 1.7% | | | | |
| 44B | Chelsea loamy fine sand, 0 to 6 percent slopes | 4.7 | 1.4% | | | | |
| 49A | Tedrow fine sand, 0 to 4 percent slopes | 1.9 | 0.6% | | | | |
| 53B | Oakville fine sand, loamy substratum, 0 to 6 percent slopes | 22.9 | 7.0% | | | | |
| 66 | Udipsamments, nearly level to gently sloping | 3.6 | 1.1% | | | | |
| 75B | Marlette-Capac loams, 1 to 6 percent slopes | 6.0 | 1.8% | | | | |
| Totals for Area of Inte | rest | 329.2 | 100.0% | | | | |

Field: W PB-W Operation: Walnutdale Farm

C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp Plan File:

Last Saved:

4/24/2013*

C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb RUSLE2 Database:

mmp_calc\Walnutdale Dairy 2013.mmp⁽⁶⁾ (6) W PB-W Calculated Profile:

CMZ 04\c.Other Local Mgt Records\mmp calc\Walnutdale Dairy 2013.mmp\(\begin{calcases} \text{(b) (6)} \end{calcases} \) Calculated Mgmt:

Field's RUSLE2 Profile

USA\Michigan\Allegan County Climate Location: Soil Survey: Allegan County, Michigan

10C OAKVILLE FINE SAND, 6 TO 18 PERCENT SLOPES\Oakville fine sand 98% Soil Type:

Slope: 400 feet Slope Length:

(Not entered -- 0% assumed) Rock Cover:

Subsurface Drainage: (No patterned drainage -- none used)

(Not selected -- rows up-and-down hill assumed) Contouring:

(Not selected -- none assumed) Strips/Barriers: (Not selected -- none assumed) Diversion/Terrace:

(Not selected -- normal burial assumed) Residue Burial:

Notes:

Field's RUSLE2 Management

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|----------------------|-------------------------------------|-------------------------|---------------------------------|
| 05/25/2008 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 405 lbs |
| 07/01/2008 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2008 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |
| 05/25/2009 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 405 lbs |
| 07/01/2009 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2009 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |
| 05/25/2010 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 405 lbs |
| 07/01/2010 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2010 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |
| 05/25/2011 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 405 lbs |
| 07/01/2011 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2011 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |

Page 1 of 2

| Michigan MARI | Manure A | pplication | n Risk Ir | ndex | | | | | | | |
|--|-----------------|----------------|---------------|-----------------|-------------------|---------------|---------------|-------------|---------------|--------------|--------------|
| Version 4 Nov 2008 | Fill in shade | d areas onl | y! | ctrl "c" to cle | ear sheet | | | | | | |
| Business Name: | Walnutda | le Dairy | | | Date | 4/24/2013 | 152 | | | | |
| Farm name: | Ned W | Ned W | (b) (6) | Bast W | Bast M | Bast E | Bast S | | | | |
| Tract Numbers: | | | | | | | | | | | |
| Section: | Dorr 07 | Dorr 07 | Dorr 12 | Dorr 14 | Dorr 14 | Dorr 14 | Dorr 23 | | | | |
| Field No: | Ned WN | Ned WS | (b) (6) | PB-W | PB-M | PB-E | PB-S | | | | |
| Acres: | 22 | 23 | 60 | 14 | 8 | 7 | 4 | | | | |
| FIELD FEATURES "INPUT" | | | | | | - | | | | | |
| I. SOIL | | | | | Mary Constitution | | | | | | |
| If drained, enter Y | | | | | | | | | | | |
| Insert Soil Series | Capac | Rimer | Capac | Oakville | Metea | Metea | Rimer | | | | |
| Soil Hydrologic Group | С | С | С | Α | В | В | С | #N/A | #N/A | #N/A | #N/A |
| Soil Management Group (SMG) | 2.5b-s | 3b | 2.5b-s | 5a | 4/2a | 4/2a | 3b | | | | |
| 3. Percent Slope | 3 | 2 | 2 | 16 | 12 | 4 | 3 | | | | |
| II. WATER QUALITY | | | | | | | | | | | |
| 4. Soil Test Phosphorus Value | 206 | 296 | 46 | | | | | | - 1 | | Land Service |
| 5. Conc. Water/Surface Inlet | S | f | f | s | s | S | s | | | | 2 4 1 2 2 3 |
| 6. Nitrogen leaching Index | m | m | m | Н | m | m | m | | | | |
| Prompt for cell above III. SURFACE COVER | Мар | Мар | Мар | Мар | m | m | Мар | #N/A | #N/A | #N/A | #N/A |
| 7. Residue/Cover Crops/Per. Cover | 70 | 70 | 70 | 70 | 40 | 50 | 50 | | | | |
| 8. Surface Water Setback | 4 | 1 | 4 | 2 | 1 | 8 | 8 | | | | |
| Vegetative Buffer Width | · | | | _ | | | | | | | |
| IV. MANURE | | | | | | | | | | | |
| 10. Manure Phosphorus Application | 58 | 58 | 94 | Ser Street | | | | | TWE WAR | | |
| 11. Manure Nitrogen Application | 70 | 70 | 145 | | 200000 | | | | | | |
| 12. Manure Application Method | SI | SI | DI | S>3 | DI | S>3 | S>3 | | | | |
| Manure Application Description | (1)DI Direct In | iect (2)SI Sur | face apply in | corp. In 2 days | (4)S<3 Surf. | apply incorp. | n 90 days (8) | S>3 Surf ar | ply left more | than 90 days | |

| Michigan MARI | Manure A | Application | n Risk Ir | ndex | | | | | | | |
|--|---------------|-------------|-----------|-----------------|-----------|-----------|--------|-----------|------------|-------------|-------|
| Version 4 Nov 2008 | Fill in shade | ed areas on | ly! | ctrl "c" to cle | ear sheet | | | | | | |
| Business Name: | Walnutda | ale Dairy | | | Date | 4/24/2013 | | | | | |
| Farm name: | Ned W | Ned W | (b) (6) | Bast W | Bast M | Bast E | Bast S | | | | |
| Tract Numbers: | | | | | | 11 | | | | | |
| Section: | | Dorr 07 | Dorr 12 | Dorr 14 | Dorr 14 | | | | | | |
| Field No: | | Ned WS | (b) (6) | PB-W | PB-M | | PB-S | | | | |
| Acres: | 22 | 23 | 60 | 14 | 8 | 7 | 4 | | | | |
| FIELD FEATURES "OUTPUT" | | | | | | | | | | | |
| | | | | | | | | | | | |
| I. SOIL | 2.5b-s | 3b | 2.5b-s | 5a | 4/2a | 4/2a | 3b | | | | |
| Soil Hydrologic Group | 4 | 4 | 4 | 1 | 2 | 2 | 4 | #N/A | #N/A | #N/A | #N/A |
| Soil Management Group | 2 | 2 | 2 | 1 | 2 | 2 | 2 | #N/A | #N/A | #N/A | #N/A |
| 3. Percent Slope | 2 | 2 | 2 | 8 | 8 | 4 | 2 | 1 | 1 | 1 | 1 |
| II. WATER QUALITY | | | | | | | | | | | |
| 4. Soil Test Phosphorus Value | 12 | 12 | 3 | | | | | | | | |
| 5. Conc. water flow or inlet discharge | 6 | 3 | 3 | 6 | 6 | 6 | 6 | FALSE | FALSE | FALSE | FALSE |
| 6. Nitrogen Leaching Index for SMG | 6 | 6 | 6 | 12 | 6 | 6 | 6 | | | | |
| III. SURFACE COVER | | | | | | | | | | | |
| 7. Residue/Cover Crop/Per. Cover | 1 | 1 | 1 | 1 | 1 | 1 | 1 | | | | |
| 8. Surface Water Setback | 4 | 1 | 4 | 2 | 1 | 8 | 8 | 0 | 0 | 0 | 0 |
| 9. Vegetative Buffer Width | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 | 12 |
| IV. MANURE | | | | | | | | | | | |
| 10. Manure "P" Application | 2 | 2 | 4 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 11. Manure "N" Application | 2 | 2 | 4 | 1 | 1 | 1 | 1 | 1 | 1 | 1 | 1 |
| 12. Manure Application Method | SI | SI | DI | S>3 | DI | S>3 | S>3 | 0 | 0 | 0 | 0 |
| Field Features Index Totals | 53 | 47 | 45 | 45 | 40 | 43 | 43 | #N/A | #N/A | #N/A | #N/A |
| | High | High | Medium | High | High | Medium | Medium | #N/A | #N/A | #N/A | #N/A |
| Total Acres by MARI risk | 0 | 0 | 71 | 67 | 138 | | | | | | |
| Category | V. low | Low | Medium | High | Total | | 0 | Total Low | and Verv I | ow risk acr | es |

Field:

W PB-W

Operation:

Walnutdale Farm

Plan File:

C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp 4/24/2013*

Last Saved:

RUSLE2 Database:

C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb

Calculated Profile:

mmp_calc\Walnutdale Dairy 2013.mmp(b) (6) W PB-W

Calculated Mgmt:

CMZ 04\c.Other Local Mgt Records\mmp_calc\Walnutdale Dairy 2013.mmp (6) (6)

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|----------------------|-------------------------------------|-------------------------|---------------------------------|
| 06/01/2012 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.2 tons | 405 lbs |
| 07/15/2012 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.2 tons | 319 lbs |
| 09/01/2012 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.7 tons | 324 lbs |
| 05/25/2013 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 449 lbs |
| 07/01/2013 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2013 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |

Note: Operations for the field's assumed previous crop (in italics) were included to initialize RUSLE2.

Field's RUSLE2 Soil Loss Estimates

| Crop Year | Starting Date mm/dd/yyyy | Ending Date mm/dd/yyyy | Soil Loss Tons/Acre | Crop(s) |
|-----------|--------------------------|------------------------|------------------------|------------------|
| 2009 | 08/11/2008 | 08/10/2009 | 1.6 | Alfalfa topdress |
| 2010 | 08/11/2009 | 08/10/2010 | 1.7 | Alfalfa topdress |
| 2011 | 08/11/2010 | 08/10/2011 | 1.7 | Alfalfa topdress |
| 2012 | 08/11/2011 | 09/01/2012 | 2.0 | Alfalfa topdress |
| 2013 | 09/02/2012 | 08/10/2013 | 1.2 | Alfalfa topdress |

RUSLE2 Version:

1.32.3.0

RUSLE2 Build Date: RUSLE2 Science Date: 20061020

Dec 17 2007

Field: M PB-M Walnutdale Farm

Plan File: C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp

Last Saved:

4/24/2013*

RUSLE2 Database: C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb

Calculated Profile: mmp calc\Walnutdale Dairy 2013.mmp\\(\begin{align*} \text{(b) (6)} \\ \text{O} \\ \text{PB-M} \end{align*}

Calculated Mgmt: CMZ 04\c.Other Local Mgt Records\mmp_calc\Walnutdale Dairy 2013.mmp\606 M PB-M

Field's RUSLE2 Profile

Climate Location: USA\Michigan\Allegan County
Soil Survey: Allegan County, Michigan

Soil Type: 27B METEA LOAMY FINE SAND, 1 TO 6 PERCENT SLOPES\Metea loamy fine sand 90%

Slope: 12% Slope Length: 300 feet

Rock Cover: (Not entered -- 0% assumed)

Subsurface Drainage: (No patterned drainage -- none used)

Contouring: (Not selected -- rows up-and-down hill assumed)

Strips/Barriers: (Not selected -- none assumed)
Diversion/Terrace: (Not selected -- none assumed)

Residue Burial: (Not selected -- normal burial assumed)

Notes:

Field's RUSLE2 Management

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|--|-------------------------------------|-------------------------|---------------------------------|
| 05/01/2008 | Planter, double disk opnr w/fluted coulter | Corn, grain | 150 bushels | |
| 10/20/2008 | Harvest, killing crop 50pct standing stubble | | | 4114 lbs |
| 05/25/2009 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1 tons | |
| 07/01/2009 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1 tons | 237 lbs |
| 08/10/2009 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.1 tons | 249 lbs |
| 05/25/2010 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1 tons | 297 lbs |
| 07/01/2010 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1 tons | 237 lbs |
| 08/10/2010 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.1 tons | 249 lbs |
| 05/25/2011 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1 tons | 297 lbs |
| 07/01/2011 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1 tons | 237 lbs |
| 08/10/2011 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.1 tons | 249 lbs |
| 05/01/2012 | Planter, double disk opnr w/fluted coulter | Corn, silage | 15 tons | 639 lbs |

Field: M PB-M Walnutdale Farm

Plan File: C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp

Last Saved:

4/24/2013*

RUSLE2 Database: C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb

Calculated Profile: mmp_calc\Walnutdale Dairy 2013.mmp\(\begin{align*} \text{(6) (6)} \\ \text{0 PB-M} \end{align*}

Calculated Mgmt: CMZ 04\c.Other Local Mgt Records\mmp_calc\Walnutdale Dairy 2013.mmp\(^{(6)}) M PB-M

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|--|-------------|-------------------------|---------------------------------|
| 09/15/2012 | Harvest, silage | | | 529 lbs |
| 05/01/2013 | Planter, double disk opnr w/fluted coulter | Corn, grain | 150 bushels | |
| 10/20/2013 | Harvest, killing crop 50pct standing stubble | | | 4114 lbs |

Note: Operations for the field's assumed previous crop (in italics) were included to initialize RUSLE2.

Field's RUSLE2 Soil Loss Estimates

| Crop Year | Starting Date mm/dd/yyyy | Ending Date mm/dd/yyyy | Soil Loss Tons/Acre | Crop(s) |
|-----------|--------------------------|---------------------------|------------------------|------------------|
| 2009 | 10/21/2008 | 08/10/2009 | 0.1 | Alfalfa topdress |
| 2010 | 08/11/2009 | 08/10/2010 | 0.3 | Alfalfa topdress |
| 2011 | 08/11/2010 | 08/10/2011 | 0.9 | Alfalfa topdress |
| 2012 | 08/11/2011 | 09/15/2012 | 1.7 | Corn silage |
| 2013 | 09/16/2012 | 10/20/2013 | 3.6 | Corn grain |

RUSLE2 Version: 1.32.3.0 RUSLE2 Build Date: Dec 17 2007 RUSLE2 Science Date: 20061020

Field: E PB-E Walnutdale Farm

Plan File: C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp

Last Saved:

4/24/2013*

RUSLE2 Database: C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb

Calculated Profile: mmp_calc\Walnutdale Dairy 2013.mmp(6) 6 E PB-E

Field's RUSLE2 Profile

Climate Location: USA\Michigan\Allegan County Soil Survey: Allegan County, Michigan

Soil Survey: Allegan County, Mich

Soil Type: 27B METEA LOAMY FINE SAND, 1 TO 6 PERCENT SLOPES\Metea loamy fine sand 90%

Slope: 4% Slope Length: 150 feet

Rock Cover: (Not entered -- 0% assumed)

Subsurface Drainage: (No patterned drainage -- none used)

Contouring: (Not selected -- rows up-and-down hill assumed)

Strips/Barriers: (Not selected -- none assumed)
Diversion/Terrace: (Not selected -- none assumed)

Residue Burial: (Not selected -- normal burial assumed)

Notes:

Field's RUSLE2 Management

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|---|----------------------|-------------------------|---------------------------------|
| 05/15/2008 | Drill or air seeder single disk openers 7-10 in spac. | Soybean, mw 7in rows | 45 bu | 1800 lbs |
| 10/10/2008 | Harvest, killing crop 20pct standing stubble | | | 1227 lbs |
| 04/28/2009 | Chisel, twisted shovel | | | |
| 04/28/2009 | Cultivator, field 6-12 in sweeps | | | |
| 05/01/2009 | planter, double disk opnr | Corn, grain | 150 bushels | |
| 10/20/2009 | Harvest, killing crop 50pct standing stubble | | | 4114 lbs |
| 10/27/2009 | Disk, tandem light finishing | | | |
| 05/15/2010 | Drill or air seeder single disk openers 7-10 in spac. | Soybean, mw 7in rows | 45 bu | |
| 10/10/2010 | Harvest, killing crop 20pct standing stubble | | | 1227 lbs |
| 05/01/2011 | Planter, double disk opnr w/fluted coulter | Corn, grain | 150 bushels | |
| 10/20/2011 | Harvest, killing crop 50pct standing stubble | | | 4114 lbs |
| 04/10/2012 | Chisel, twisted shovel | | | |

Field:

E PB-E Walnutdale Farm

Operation: Plan File:

C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp 4/24/2013*

Last Saved:

RUSLE2 Database:

C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb

Calculated Profile:

mmp_calc\Walnutdale Dairy 2013.mmp\(\begin{cases} \begin{

CMZ 04\c.Other Local Mgt Records\mmp_calc\Walnutdale Dairy 2013.mmp 6 6 E PB-E Calculated Mgmt:

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|----------------------------------|---|-------------------------|---------------------------------|
| 04/15/2012 | Cultivator, field 6-12 in sweeps | | | |
| 04/15/2012 | Drill or airseeder, double disk | Alfalfa, spring seed | 1 Tons | |
| 07/15/2012 | Harvest, hay, legume | Alfalfa, spring seed regrowth after cutting | 1 tons | 270 lbs |
| 09/01/2012 | Harvest, hay, legume | Alfalfa, spring seed senes to y2 regrowth | 2.5 tons | 270 lbs |
| 05/25/2013 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 661 lbs |
| 07/01/2013 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2013 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |

Note: Operations for the field's assumed previous crop (in italics) were included to initialize RUSLE2.

Field's RUSLE2 Soil Loss Estimates

| Crop Year | Starting Date mm/dd/yyyy | Ending Date mm/dd/yyyy | Soil Loss Tons/Acre | Crop(s) |
|-----------|--------------------------|------------------------|------------------------|------------------|
| 2009 | 10/11/2008 | 10/20/2009 | 1.6 | Corn grain |
| 2010 | 10/21/2009 | 10/10/2010 | 0.7 | Soybean |
| 2011 | 10/11/2010 | 10/20/2011 | 1.0 | Corn grain |
| 2012 | 10/21/2011 | 09/01/2012 | 0.6 | Alfalfa seeding |
| 2013 | 09/02/2012 | 08/10/2013 | 0.5 | Alfalfa topdress |

RUSLE2 Version: RUSLE2 Build Date:

1.32.3.0 Dec 17 2007 RUSLE2 Science Date: 20061020

S PB-S Field: Walnutdale Farm Operation:

C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp Plan File:

Last Saved:

4/24/2013*

RUSLE2 Database: C:\Program Files (x86)\USDA\Rusle2\Import\BASE NRCS MOSES 040110.gdb

mmp calc\Walnutdale Dairy 2013.mmp\(\begin{calcor} Calculated Profile:

CMZ 04\c. Other Local Mgt Records\mmp calc\Walnutdale Dairy 2013.mmp\(\begin{calcrage}{c} \text{D} \text{(6)} \text{(6)} \end{calcrage} \) Calculated Mgmt:

Field's RUSLE2 Profile

Climate Location: USA\Michigan\Allegan County Soil Survey: Allegan County, Michigan

Soil Type:

28A RIMER LOAMY SAND, 0 TO 4 PERCENT SLOPES\Rimer loamy sand 90%

Slope: 3% Slope Length: 150 feet

Rock Cover: (Not entered -- 0% assumed)

Subsurface Drainage: (No patterned drainage -- none used)

(Not selected -- rows up-and-down hill assumed) Contouring:

Strips/Barriers: (Not selected -- none assumed) Diversion/Terrace: (Not selected -- none assumed)

Residue Burial: (Not selected -- normal burial assumed)

Notes:

Field's RUSI F2 Management

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed | |
|--------------------|---|-------------------------------------|-------------------------|---------------------------------|--|
| 05/25/2008 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 405 lbs | |
| 07/01/2008 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs | |
| 08/10/2008 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs | |
| 04/24/2009 | Disk, tandem secondary op. | | | 2378 lbs | |
| 04/28/2009 | Cultivator, field 6-12 in sweeps | | | | |
| 05/01/2009 | planter, double disk opnr | Corn, grain | 125 bushels | | |
| 10/20/2009 | Harvest, killing crop 50pct standing stubble | | | 3471 lbs | |
| 05/05/2010 | Disk, tandem secondary op. | | | | |
| 05/15/2010 | Cultivator, field 6-12 in sweeps | | | | |
| 05/15/2010 | Drill or airseeder, double disk | Soybean, mw 7in rows | 40 bu | | |
| 09/20/2010 | Harvest, killing crop 20pct standing stubble | | | 1346 lbs | |
| 09/25/2010 | Drill or air seeder single disk openers 7-10 in spac. | Rye, cereal | 35 bushels | | |

Field: 5 PB-S Walnutdale Farm

Plan File: C:\Local Cloud\Shared\Manure Plan Documents\Walnutdale Dairy CNMP Data\Walnutdale Dairy 2013.mmp

Last Saved:

4/24/2013*

RUSLE2 Database: C:\Program Files (x86)\USDA\Rusle2\Import\BASE_NRCS_MOSES_040110.gdb

Calculated Profile: mmp_calc\Walnutdale Dairy 2013.mmp (6) S PB-S

Calculated Mgmt: CMZ 04\c.Other Local Mgt Records\mmp_calc\Walnutdale Dairy 2013.mmp⁽⁶⁾ S PB-S

| Date mm/dd/yyyy | Operation | Vegetation | Yield or Cover Material | Cover Material Added/Removed |
|--------------------|--|--|-------------------------|---------------------------------|
| 07/20/2011 | Harvest, killing crop 50pct standing stubble | | | 1491 lbs |
| 08/15/2011 | Drill or airseeder, double disk | Alfalfa, fall seed senes to yr2 regrowth | 0.7 tons | |
| 06/01/2012 | Harvest, hay, legume | Alfalfa, yr2 regrowth after cutting | 0.7 tons | 179 lbs |
| 07/15/2012 | Harvest, hay, legume | Alfalfa, yr2 regrowth after cutting | 0.7 tons | 186 lbs |
| 09/01/2012 | Harvest, hay, legume | Alfalfa, yr2 regrowth after cutting | 2.5 tons | 189 lbs |
| 05/25/2013 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 675 lbs |
| 07/01/2013 | Harvest, hay, legume | Alfalfa, yr4 regrowth after cutting | 1.3 tons | 308 lbs |
| 08/10/2013 | Harvest, hay, legume | Alfalfa, yr4 senes to yr5 regrowth | 1.5 tons | 324 lbs |

Note: Operations for the field's assumed previous crop (in italics) were included to initialize RUSLE2.

Field's RUSLE2 Soil Loss Estimates

| Crop Year | Starting Date mm/dd/yyyy | Ending Date mm/dd/yyyy | Soil Loss Tons/Acre | Crop(s) |
|-----------|--------------------------|------------------------|------------------------|------------------|
| 2009 | 08/11/2008 | 10/20/2009 | 0.9 | Corn grain |
| 2010 | 10/21/2009 | 09/20/2010 | 0.6 | Soybean |
| 2011 | 09/21/2010 | 07/20/2011 | 0.4 | Rye grain |
| 2012 | 07/21/2011 | 09/01/2012 | 0.7 | Alfalfa seeding |
| 2013 | 09/02/2012 | 08/10/2013 | 0.2 | Alfalfa topdress |

RUSLE2 Version: 1.32.3.0 RUSLE2 Build Date: Dec 17 2007 RUSLE2 Science Date: 20061020

| | | Win | d Erosion W | orksheet | | | | |
|--------------|------------------|-----------------|--------------------|-------------|---------------|-------------|-------------------|--|
| | base | d on Wind Eros | | | ection 1 F | OTG | | |
| c | lient: | | d #: | Date: | | | County | |
| | nutdale | (b) (6) | PB-W | | /2013 | | llegan | |
| | | Soil "I" Value | e - refer to Sect | | | | 0 | |
| | Na | me | # | I value | Т | | | |
| Call Tuna 1 | Oak | n illa | 100 | 250 | | 1 | | |
| Soil Type 1: | | ville | 10C | 250 | 5 | - | | |
| Soil Type 2: | | lette | 14C | 56 | 5 | | | |
| | Soil Roug | hness (Ridge) \ | | fer to Tabl | | n 1 FOTG | | |
| | 3 | Tillage used | for Krd | | Krd Value | | | |
| Present | | none | | | 1.0 | | | |
| Planned | | none | | | 1 | | | |
| | | | or - refer to tabl | | 1 FOTG | | | |
| County: | Alle | gan | Climatic F | | 8 | | | |
| | | | ngth of Unshelt | ered Distai | nce | | | |
| | | Windbreak adju | | | culated "L" (| | Adjusted "L" | |
| Soil Type 1: | Measured "L" | Height | Туре: | Angle | Adj factor | Field width | value (Ft) | |
| Present | 670 | | fenceline | | | | 470 | |
| Planned | 670 | | fenceline | | | | 470 | |
| | | Windbreak adju | | | culated "L" (| | Adjusted "L" | |
| Soil Type 2: | Measured "L" | Height | Туре: | Angle | Adj factor | Field width | value (Ft) | |
| Present | 340 | | fenceline | | | | 140 | |
| Planned | 340 | | fenceline | | | | 140 | |
| | "V" - Vegitative | Factor (Small (| | | | | | |
| | | | % residue | Lbs | Manure | SGe (Fig. a | -1 to c-2.) Table | |
| | Present Crop | Residue | Cover | residue | rate | | 1 | |
| Soil Type 1: | Alfalfa | Alfalfa | 60% | | 0 | | 2700 | |
| Soil Type 2: | Alfalfa | Alfalfa | 60% | | 0 | | 2700 | |
| | | | % residue | Lbs | Manure | SGe (Fig. a | -1 to c-2.) Table | |
| | Planned Crop | Residue | Cover | residue | rate | | 2 | |
| Soil Type 1: | Alfalfa | Alfalfa | 60% | | 0 | 2700 | | |
| Soil Type 2: | Alfalfa | Alfalfa | 60% | | 0 | | 2700 | |
| | "E" Estimated A | | | | | | | |
| | Present Crop | Soil Loss | | Soil Loss | Average | Soil Loss | Tolerable (T/ac) | |
| Soil Type 1: | Alfalfa | | Alfalfa | 0 | | 0 | 5 | |
| Soil Type 2: | Alfalfa | 0 | Alfalfa | 0 | | 0 | 5 | |

| | | Win | d Erosion We | orksheet | | | |
|--------------|------------------|-----------------|--------------------|-------------|---------------|-------------|--------------------|
| | base | d on Wind Eros | | | ection 1 F | OTG | |
| С | lient: | | d #: | | ite: | | ounty |
| | Inutdale | (b) (6) | PB-M | 4/24 | /2013 | | llegan |
| | | Soil "I" Value | e - refer to Sect | | | | 0 |
| i | Na | me | # | I value | Т | | |
| Call Time 4 | Dell | ville | 48 | 134 | 4 | I | |
| Soil Type 1: | Bell | VIIIe | | | 4 | - | |
| Soil Type 2: | | ville | 10C | 250 | 5 | | |
| | Soil Roug | hness (Ridge) \ | | fer to Tabl | | n 1 FOTG | |
| | | Tillage used | for Krd | | Krd Value | | |
| Present | | None | | | 1.0 | | |
| Planned | | None | | | 1 | | |
| | | | or - refer to tabl | | 1 FOTG | | |
| County: | Alle | gan | Climatic F | | 8 | | |
| | , | | ngth of Unshelt | ered Distai | тсе | | |
| | | Windbreak adju | | | culated "L" (| | Adjusted "L" |
| Soil Type 1: | Measured "L" | Height | Туре: | Angle | Adj factor | Field width | value (Ft) |
| Present | 670 | | fenceline | | | | 470 |
| Planned | 670 | | fenceline | | | | 470 |
| | | Windbreak adju | | | culated "L" (| | Adjusted "L" |
| Soil Type 2: | Measured "L" | Height | Туре: | Angle | Adj factor | Field width | value (Ft) |
| Present | 600 | | fenceline | | | | 400 |
| Planned | 600 | | fenceline | | | | 400 |
| | "V" - Vegitative | Factor (Small | | | | | |
| | | | % residue | Lbs | Manure | SGe (Fig. a | ı-1 to c-2.) Table |
| | Present Crop | Residue | Cover | residue | rate | | 1 |
| Soil Type 1: | Silage Corn | Silage Corn | | | 9000 Gal | | 700 |
| Soil Type 2: | Silage Corn | Silage Corn | | | 9000 Gal | | 700 |
| | | | % residue | Lbs | Manure | SGe (Fig. a | -1 to c-2.) Table |
| | Planned Crop | Residue | Cover | residue | rate | | 2 |
| Soil Type 1: | | Grain Corn | 60% | | | | |
| Soil Type 2: | Grain Corn | Grain Corn | 60% | | | | |
| | "E" Estimated A | | | | | | |
| | Present Crop | Soil Loss | | Soil Loss | Average | Soil Loss | Tolerable (T/ac |
| Soil Type 1: | Silage Corn | | Grain Corn | 0.2 | | 0.45 | 4 |
| Soil Type 2: | Silage Corn | 3.4 | Grain Corn | 1.3 | | 2.35 | 5 |

| | | Win | d Erosion We | orksheet | | | |
|-------------------------|------------------|--------------------------|-------------------------------|---------------|--------------------|--------------------|---------------------------------------|
| | base | d on Wind Eros | ion Worksheet | found in S | ection 1 F | OTG | |
| С | lient: | Fiel | ld #: | Da | ite: | C | ounty |
| | Inutdale | (b) (6) | PB-E | 4/24/ | /2013 | | llegan |
| | | Soil "I" Value | e - refer to Sect | ion II of the | FOTG | | |
| | Na | me | # | I value | Т | | |
| Soil Type 1: | Me | tea | 27B | 134 | 4 |] | |
| | | | | | | 1 | |
| Soil Type 2: | 0 11 0 | (D: 1)) | (1 (1) | f (T) I | F 0 1 | 4 5070 | |
| | Soil Roug | hness (Ridge) \ | | fer to Tabl | | n 1 FOIG | |
| Б., | | Tillage used | for Kra | | Krd Value | - | |
| Present | | none | | | 1.0 | | |
| Planned | | none | or - refer to tabl | - 0 04: | 4 5050 | | |
| 0 | ۸۱۱۵ | | | | | | |
| County: | Alle | gan | Climatic F ngth of Unshelt | | 8 | | |
| | T | | | Or cold | culated "L" (| Table 4) | Adjusted "L" |
| Cail Type 1: | Measured "L" | Windbreak adju Height | | | | Field width | |
| Soil Type 1: Present | 1000 | | Type: fenceline | Angle | Auj lactor | Field Width | value (Ft) 800 |
| Planned | 1000 | | fenceline | | | | 800 |
| Planned | 1000 | Windbreak adju | | Or cold | L culated "L" (| Table 4) | Adjusted "L" |
| Soil Type 2: | Measured "L" | Height | Type: | Angle | | Field width | value (Ft) |
| Present | INICASUICU L | rieigiit | Гуре. | Angle | Auj lactor | i iela wiatii | value (1 t) |
| Planned | | | | | | | 0 |
| 1 latified | "V" - Vegitative | Factor (Small) | l Grain equivaler | t) for each | cron in ro | l tational neri | |
| | VVegitative | Taotor (oman | % residue | Lbs | Manure | | -1 to c-2.) Table |
| | Present Crop | Residue | Cover | residue | rate | 000 (1 19. 0 | 1 |
| Soil Type 1: | Alfalfa | Alfalfa | 40% | | | | · · · · · · · · · · · · · · · · · · · |
| Soil Type 2: | / wand | 7 ilialia | 1070 | 1000 | | | |
| 3011 Type 2. | | | % residue | Lbs | Manure | SGe (Fig. a | -1 to c-2.) Table |
| | Planned Crop | Residue | Cover | residue | rate | 2 | |
| Soil Type 1: | Alfalfa | Alfalfa | 60% | | 0 | | |
| Soil Type 2: | | | | | | | |
| | "E" Estimated A | nnual Soil Loss | s by wind Erosi | on (from "I | E" tables S | ection 1 FO | TG) |
| | Present Crop | Soil Loss | | Soil Loss | _ | Soil Loss | Tolerable (T/ac) |
| Soil Type 1: | Alfalfa | | Alfalfa | 0.1 | 3- | 0.65 | 4 |
| Soil Type 2: | | | | | | | |

| | | Win | d Erosion We | orksheet | | | | |
|--------------|------------------|-----------------|--------------------|---------------|---------------|-------------|-------------------|--|
| | base | d on Wind Eros | ion Worksheet | found in S | ection 1 Fo | OTG | | |
| CI | lient: | Fiel | d #: | Da | ite: | C | County | |
| Wal | nutdale | (b) (6) | PB-S | 4/24 | /2013 | Α | llegan | |
| | | Soil "I" Value | e - refer to Sect | ion II of the | FOTG | | | |
| | Na | me | # | I value | Т | | | |
| Soil Type 1: | Pipes | stone | 26A | 220 | 5 | | | |
| Soil Type 2: | | ville | 53B | 250 | 4 | | | |
| | Soil Roug | hness (Ridge) \ | /alue (Krd) - *re | fer to Tabl | e 5, Sectio | n 1 FOTG | | |
| | | Tillage used | | | Krd Value | | | |
| Present | | none | | | 1.0 | | | |
| Planned | | none | | | 1 | | | |
| | | Climatic Facto | or - refer to tabl | e 2 Section | 1 FOTG | | | |
| County: | Alle | gan | Climatic F | | 8 | | | |
| | | | ngth of Unshelt | | | | | |
| | | Windbreak adju | stments: | Or cald | culated "L" (| Table 4) | Adjusted "L" | |
| Soil Type 1: | Measured "L" | Height | Туре: | Angle | Adj factor | Field width | value (Ft) | |
| Present | 670 | | fenceline | | | | 370 | |
| Planned | 670 | | fenceline | | | | 370 | |
| | | Windbreak adju | | | culated "L" (| | Adjusted "L" | |
| Soil Type 2: | Measured "L" | Height | Туре: | Angle | Adj factor | Field width | value (Ft) | |
| Present | 820 | | fenceline | | | | 620 | |
| Planned | 820 | | fenceline | | | | 620 | |
| | "V" - Vegitative | Factor (Small | | | | | | |
| | | | % residue | Lbs | Manure | SGe (Fig. a | -1 to c-2.) Table | |
| | Present Crop | Residue | Cover | residue | rate | | 1 | |
| Soil Type 1: | Alfalfa | Alfalfa | 40% | 10 (10 (10) | | | 2700 | |
| Soil Type 2: | Alfalfa | Alfalfa | 40% | | | | 2700 | |
| | | | % residue | Lbs | Manure | SGe (Fig. a | -1 to c-2.) Table | |
| | Planned Crop | Residue | Cover | residue | rate | | 2 | |
| Soil Type 1: | Alfalfa | Alfalfa | 60% | | | | 2700 | |
| Soil Type 2: | Alfalfa | Alfalfa | 60% | | | | 2700 | |
| ' | "E" Estimated A | | | | | | | |
| | Present Crop | Soil Loss | Planned Crop | Soil Loss | | Soil Loss | Tolerable (T/ac) | |
| Soil Type 1: | Alfalfa | | Alfalfa | 0.3 | | 0.4 | 5 | |
| Soil Type 2: | Alfalfa | 3.2 | Alfalfa | 1.2 | | 2.2 | 4 | |